



# Protocol Booklet

<b>Product Code(s)</b>	HB9578
<b>Product Name(s)</b>	ING-2 AM
<b>Purpose</b>	Measurement of intracellular Na <sup>+</sup> in cultured cells

**Please note:** This product is for RESEARCH USE ONLY and is not intended for therapeutic or diagnostic use. Not for human or veterinary use



## Contents

<b>Product Overview</b> .....	3
<b>Components &amp; Storage</b> .....	3
<b>Protocol</b> .....	3
<b>Recipes</b> .....	4
ING-2 Loading Solution .....	4
HEPES-buffered Hank's Balanced Salt Solution (Assay Buffer) .....	4
<b>Guidelines, precautions, troubleshooting</b> .....	4
<b>Contact</b> .....	4
For customers in the UK, Europe and Rest of the World .....	4
For customers in the USA, Canada and South America .....	4



## Product Overview

ING-2 AM is a membrane permeable, yellow-green fluorescent (Excitation 525nm, Emission 545nm), intracellular sodium ( $\text{Na}^+$ ) indicator ( $K_d = 20\text{mM}$ ). It has improved cellular loading and significantly higher brightness than SBF1. Suitable for high-throughput screening applications targeting  $\text{Na}^+$  channels, and non-selective monovalent cation channels due to its spectral properties and large dynamic range. Also compatible with fluorescence microscopy using common fluorescein, GFP or more ideally YFP filters.

## Components & Storage

ING-2 AM is provided as:

SKU	Component	Quantity	Storage Temperature
HB9578	ING-2 AM	500 $\mu\text{g}$	-20°C

This protocol additionally requires:

Component	Quantity	Storage Temperature
DMSO	25 $\mu\text{l}$	RT
Pluronic F-127	10mg	4°C
Probenecid	7.7mg	4°C
Assay Buffer (HEPES-buffered Hank's Balanced Salt Solution (pH = 7.3)*)	10ml	RT

\* Please see recipe at the end of this protocol book.

## Protocol

The following protocol provides general guidelines for using ING-2 AM to measure intracellular sodium in cultured cells. All loading conditions (dye concentration, temperature, and time) should be optimized for your specific assay, application, and instrumentation.

1. Culture cells following standard protocols to approximately 80-100% confluence.
2. Prepare the loading solution freshly following the below table, vortex well and use within 2 hours.
3. Remove the cell culture medium, briefly wash in plain media (without serum), then add dye loading solution. Recommend volumes are:
  - a. 35mm dish / 6-well plate - 1.5 mL/well,
  - b. 96 well plate - 100  $\mu\text{L}$ /well,
  - c. 384 well plate - 20  $\mu\text{L}$ /well,
4. Incubate in a cell culture incubator at 37°C for 60 minutes.
5. Read fluorescence using a plate reader (Excitation: 525nm, Emission 545nm) or image using a fluorescence microscope using a compatible filter set (e.g. FITC, GFP, YFP).



## Recipes

### ING-2 Loading Solution

Component	Concentration	Quantity	Notes
ING-2 AM	4.6µM	50µg	Dissolve in DMSO then aliquot and store any unused dye at -20°C
Assay Buffer	1X	10ml	Normally HEPES buffered HBSS but other buffers have been also successfully used.
Pluronic F-127	0.1%	10mg	Surfactant that helps the dissolution of dye therefore ensuring even dye distribution and cellular loading.
Probenecid	2.7mM	7.7mg	Anion transport inhibitor that improves intracellular dye retention. Not required for all cell types, it is recommended in most cases to optimize assay performance.

**Please note:** Combine components then vortex thoroughly. Use within 2 hours of creation. Do not freeze.

### HEPES-buffered Hank's Balanced Salt Solution (Assay Buffer)

Component	MW (g/mol)	g/L	Concentration (mM)
Calcium Chloride	110.98	0.14	1.26
Magnesium Chloride Hexahydrate	203.30	0.1	0.49
Magnesium Sulfate Heptahydrate	246.47	0.1	0.41
Potassium Chloride	74.55	0.4	5.33
Potassium Phosphate Monobasic	136.09	0.06	0.44
Sodium Bicarbonate	84.01	0.35	4.17
Sodium Chloride	58.44	8	138.00
Sodium Phosphate Dibasic	141.96	0.048	0.34
D-Glucose (Dextrose)	180.16	1	5.56
HEPES	238.30	4.76	20.00

**Please note:** Add all components to dH<sub>2</sub>O, mix well then adjust to pH 7.3

## Guidelines, precautions, troubleshooting

Please contact our technical support team at [technicalhelp@hellowbio.com](mailto:technicalhelp@hellowbio.com) for advice on how to resolve any problems encountered when using this product. Observe safe laboratory practice and consult the safety datasheet. Please see the datasheet on our website for general guidelines, precautions, limitations on the use of the product.

## Contact

### For customers in the UK, Europe and Rest of the World

Customer Care [customercare@hellowbio.com](mailto:customercare@hellowbio.com)

Technical support [technicalhelp@hellowbio.com](mailto:technicalhelp@hellowbio.com)

By telephone: +44(0)117 318 0505

By fax: +44(0)117 981 1601

Opening hours: 8.30 am - 5.00 pm GMT weekdays

### For customers in the USA, Canada and South America

Customer Care [customercare-usa@hellowbio.com](mailto:customercare-usa@hellowbio.com)

Technical support [technicalhelp@hellowbio.com](mailto:technicalhelp@hellowbio.com)

By telephone: +1-609-683-7500

By fax: +1-609-228-4994

Opening hours: 9.00 am - 5.00 pm EST weekdays